

INDEX.

- Acari found in moles' nests, A. D. Michael on, 269.
- Acolophus, 58, 66; debilis, 60, 66.
- Ædemonus (Erichsonii), 335.
- Æluroidea, brain of, 12, 24.
- Ælurus, brain of, 12.
- African Curculionidæ, new genera and species of, F. P. Pascoe on, 318.
- Aglaia acuminata, 304.
- Aglaophenia, 150; acutidentata, 151, 160; chalarocarpa, 138, 150, 160; dolichocarpa, 152, 161; latecarinata, 151, 152, 161; myriophyllum, 155; perforata, 150, 160; perpusilla, 152.
- Allman, Prof. G. J., Australian, Cape, and other Hydroida, from the Collection of Miss H. Gatty, 132.
- Alona acanthoceroideus, 295.
- Amblypneustes ovum, 179, 193.
- Ambulacra, of *Aspidodiadema microtuberculatum*, 95, 113; of *Astropyga pulvinata*, 95, 110; of *A. radiata*, 95, 107; of *Centrostephanus*, 95, 110; of *Diadema setosum*, 95, 96; of *Echinothrix calamaris*, 95, 106; of *E. Desori*, 95, 101; of *Micropyga tuberculata*, 95, 110.
- of recent *Diadematiidæ*, on anatomy of, Prof. P. M. Duncan, 95.
- Ambulacral compound tubercle-bearing plate, 102.
- Anatomy of ambulacra of recent *Diadematiidæ*, by Prof. P. M. Duncan, 95.
- of *Sphærotherium*, by G. C. Bourne, 161.
- Andrena fulva*, 87.
- Antennæ of Honey-Bee (Briant), 84.
- Antennary organ and nerve of *Sphærotherium*, 173.
- Antibothrus*, 124; carinatus, 124.
- Antroderus*, 126; costatus, 126, 131.
- Arbacia nigra*, 53, 57; punctulata, 49, 57; pustulosa, 49, 57; stellata, 49, 57.
- , ambulacra of *A. nigra*, 53; structure of vertical sutures of interradia, 53; structure of ambulacral plates, 48; test of, 25.
- Arbaciadæ*, apical sutures, 38; interradial plates and sutures, 47; obliquity of interradial plates, 38; test of Fam. of, Prof. Duncan and W. P. Sladen on, 25; radial plates of, 48.
- Arctietis*, brain of, 7.
- Arctoidea*, brain of, 10, 20, 24.
- Ascaris simplex*, 176; delphini, 176.
- Aspidodiadema microtuberculatum*, 95, 113.
- Astropyga radiata*, 95, 106, 110, 114; pulvinata, 95, 106, 110.
- Attelabinæ* (African), 319.
- Attelabus chrysideus*, 319, 330; costipennis, 330.
- Attheyella cingalensis*, 296; spinosa, 296.
- Aulonosoma tenebrioides*, 122.
- Australian, Cape, and other Hydroida, from the Collection of Miss H. Gatty, by Prof. G. J. Allman, 132.
- Bairdia tenera*, 304.
- Balaninæ* (African), 319.
- Balaninus*, 332; brevirostris, 319, 331; villosus, 331.
- Baly, J. S., Colombian species of genus *Diabrotica*.—Part I., 213; Part II., 230.
- Bassaricyon*, brain of, 13.
- Bassaris*, brain of, 12.
- Bell, Prof. F. J., new species of *Minyad* (*Minyas torpedo*) from N.W. Australia, 114.
- Binturong*, brain of, 7.
- Bittium abruptum*, 14.
- Bothrideres*, 75, 124, 127; contractus, 128; bituberculatus, 128.
- Bothriderini*, 60.
- Bourne, G. C., on anatomy of *Sphærotherium*, 161.
- Bousfield, E. C., on *Slavina* and *Ophidionais*, 264.
- Brachyderinæ* (African), 318.
- Brady, Dr. G. S., Notes on Entomostrea collected by Mr. A. Haly in Ceylon, 293.
- Brains of Carnivora, Prof. Mivart on, 1.

- Briant, T. J., on antennæ of honey-bee, 84.
 Bythocythere retusa, 315.
- Calandrinæ (African), 319.
 Calanidæ (Ceylon), 296.
 Calytops granosa, 327.
 Campanularia carduella, 132, 158.
 Canis Azaræ, brain of, 3, 4; microtis, brain of, 3, 4; procyonoides, brain of, 3.
 Capybara, 174, 176.
 Carnivora, St. George Mivart on cerebral convolutions of, 1.
 Cat, brain of, 2, 5.
 Catamonus melancholicus, 325; suffusus, 319, 324.
 Caulinia, 141.
 Cautomus, 58, 60; hystriculus, 60, 82, 84.
 Cavia Cobaya, 174.
 Centrostephanus, ambulacra of, 110.
 Cercoleptes, brain of, 13.
 Cerebral convolutions of Carnivora, St. George Mivart on, 1.
 Cerithiopsides from N. Atlantic and Madeira, by Rev. R. Boog Watson, 89.
 Cerithiopsis, 89; angustissima, 94; atalaya, 90, 94, 95; Barleei, 90, 91, 92, 95; carinata, 89; Clarkii, 90, 92, 95; costulata, 89, 91, 95; diadema, 90, 92, 95; Emersonii, 89; fayalensis, 90, 91, 92, 95; gemmulifera, 89; Jeffreyi, 89, 90; Metaxæ, 89, 90, 94, 95; pulchella, 89, 90, 95; rugulosa, 94; terebralis, 89; tiara, 90, 92, 95; trilineatum, 89; tubercularis, 89, 90, 95; Whiteavesii, 89.
 Cerithium metula, 89; trilineatum, 89.
 Cerotoma Deyrollii, 248.
 Cerylon, 59, 60; angustatum, 80; crasipes, n. sp., 60, 80, 81; curticolle, n. sp., 60, 81; deplanatam, 80; gracilipes, 130; minimum, 60, 81; orientale, 131; pusillum, 131; tibiale, 130, 131; quadricolle, 131.
 Cerylonini, 60.
 Ceylon Entomostraca, G. S. Brady on, 293.
 Chætogaster, 265.
 Cheetah, Prof. Owen on brain of, 1.
 Chilopoda, 166, 167, 171.
 Chlamydotheca, 293, 301; subglobosa, 300, 301.
 Cicones, 58, 60, 121; bitomoides, 60, 69, 121; coloratus, 121; oblongus, 60, 68; oculatus, 60, 67; minimus, 60, 69, 121; minutus, 121; niveus, 60, 68, 69, 121; variegatus, 68.
- Cidaridæ, 180, 181, 182, 186, 207, 208, 209; perignathic girdle of, 179, 182.
 Cidaris, 180, 184, 185, 193, 199, 202, 209.
 Civet, brain of, 5.
 Cladocera, 294.
 Cloelia fasciata, 178.
 Clypeaster humilis, 179, 203, 205, 212; rosaceus, 179, 203, 204, 205, 212; scutiformis, 203; subdepressus, 203.
 Clypeastridæ, 179, 203, 207, 209; perignathic girdle in, 179, 203.
 Clypeastroids, 181.
 Cobbold, Dr. T. Spencer, Notes on Parasites collected by the late Chas. Darwin, 174; Strongylus Axei (Cobb.), preceded by remarks on its affinities, 259.
 Cœlopleuri, ambulacra between large tubercles of ambitus and radial plates, 43; ambulacral region at ambitus, 40; peristomial region of ambulacra, 46; test recent sp. of, 39.
 Cœlopleurus, 27, 98, 106, 110, 113; equis, 27, 57; Forbesi, 27, 37; Mail-lardi, 39, 57; Pratti, 27, 56; sindensis, 27, 36, 57, 98.
 —, aboral demi-plate of, 30; adoral demi-plate of, 31; ambulacra of, 27; ambulacra near peristome, 32; ambulacra near radial plates, 33; apical sutures of, 38; central or primary plate of, 31; obliquity of interradsial plates of, 38; pits along median suture, 33; radial plates of, 37; region of the ambitus in, 29; test, fossil sp. of, 27.
 Colobicus, 58; emarginatus, var., 60, 66; granulatus, 60, 65, 121; indicus, 121; rugosulus, 120.
 Coluber, 177; flaviventris, 178.
 Colydiidæ, 58, 59, 60, 127; from Ceylon, by D. Sharp, 117; from Japan, by D. Sharp, 58.
 Colydiina, 60.
 Colydiini, D. Sharp on, 60.
 Comparisons between tests of Cœlopleurus and Arbacia, 56.
 Convolutions, cerebral, of Carnivora, Prof. St. G. Mivart on, 1.
 Copepoda (Ceylon), 296.
 Copulatory appendages in Sphærothetherium, 164.
 Coxelus unicolor, 119.
 Crocuta, brain of, 9, 10.
 Crossarchus, brain of, 8.
 Crustacea, 262.
 Cryptoprocta, brain of, 9, 10.
 Cryptorhynchinæ (African), 319.
 Curculionidæ, new African genera and species of, F. P. Pascoe on, 318.

- Cyclopidae, 296.
 Cyclops, sp., from Ceylon, 296.
 Cylindromicrus, 58, 60; gracilis, 60, 73, 84.
 Cynogale, brain of, 7.
 Cynoidea, Prof. Mivart on cerebral convolutions of, 3, 24.
 Cypridea, defined by Prof. Rupert Jones, finote, 300; Bosquet's genus Cypridea, 301.
 Cyprididae (Ceylon), 297.
 Cypridopsis globosa, 303; marmorata, 303.
 Cyprinotus, 293, 301; cingalensis, 302.
 Cypris, affinis, 299; brasiliensis, 301; (Chlamydotheca) Azteca, 301; cylindrica, 293, 301; cylindrica, var. major, 297; furfuracea, 299; Hayli, 299; luxata, 298; Malcolmsoni, 293; monilifera, 298; obliqua, 299; purpurascens, 298; subglobosa, 293, 300; tenuicauda, 299; tessellata, 299.
 Cystophora, brain of, 21.
 Cythere Andei, 310; bimammillata, 309; cancellata, 306; coralloides, 306; Darwini, 308; fabacea, 305; Goujoni, 308; Hodgii, 310; iniqua (= Cytherura bataviana), 310; lactea, 309; laqueata, 308; melobesioides, 309; nodulifera, 309; Normaniana, 308; papuensis, 309; rectangularis, 310; Ruperti, 306; Stimpsoni, 308; subcuneata, 306; trunculata, 305.
 Cytheridae of Ceylon, 305.
 Cytheridea orientalis, 311; pusilla, 311.
 Cytherura bataviana (= Cythere iniqua), 310, 311.
 Daphnidae, 294.
 Daresté, M., memoir on convolutions of mammalian brain, 1.
 Dastarcus, 59; longulus, 60, 76, 84; porosus, 76, 128.
 Delphinus Forsteri and D. phocæna, parasites in, 177.
 Deretaphrini, 60.
 Dermaleichi, coitus of, 273.
 Dermatodes metallescens, 319, 330.
 Desmidophorus encaustus, 319, 333; fascicularis, 332; penicillatus, 332; Satanas, 319, 332, 336; ursus, 333.
 Desmoscyphus orifissus, 143, 159; unguiculata, 144, 160.
 Desmosticha, 203.
 Diabrotica, Colombian Species of, by J. S. Baly, 213, 230.
 —, 213; abbreviata, 238; abruptum, 259; adonis, 239; æneipennis, 234; alternata, 244; amabilis, 222; arcuata, 218; beata, 244, 249, 251; bella, 246, 247; biannularis, 219, 220; bipustulata, 221; bivittata, 233; bivittaticollis, 257, 258; Butleri, 251, 252; Chapuisi, 227; Chevrolati, 256; chrysopleura, 215; cincella, 238; circulata, 243; clypeata, 249, 251; coccinea, 234; consentanea, 214; corusca, 230; coryphæa, 233; decempuncta, 220, 221; deliciosa, 251; Deyrollii, 241; Dysoni, 217; elegantula, 213; fenestrata, 250; flavo-cincta, 235; flavo-limbata, 234; flavo-marginata, 234; formosa, 226; fulvogignata, 229; fusco-maculata, 258, 259; Gemmingeri, 225; gloriosa, 239; gratiosa, 215; Haroldi, 219, 220; Hebe, 241; hexaspilota, 228; histrionica, 215, 216; imitans, 251; inæqualis, 219; incerta, 236; inconstans, 224, 225; innuba, 230; insignita, 256; instabilis, 258; intermedia, 252; Jacobyi, 259; Jekelii, 245, 246; jucunda, 247; Kirschi, 231, 232; Klugii, 258; labiata, 223; Lacordairei, 259; læta, 248, 249; lætabilis, 257; Lebasii, 221; leucospila, 253; lugubris, 253; mediiovittata, 237; mimula, 245, 246; mutabilis, 225, 226; nigriceps, 235; nigroguttata, 243; nigrolimbata, 218; nigronotata, 216; nigrovittulata, 242; ornatula, 224, 258; piceo-lineata, 243; placida, 220; porracea, 224; posticata, 252; propinqua, 255; puella, 238; pulchra, 239; puncticollis, 234; regalis, 213; robusta, 255, 256; Sallei, 227; separata, 232; serraticornis, 255; sexplagiata, 257; similata, 235, 237; simulans, 222; spectanda, 241; spilopectera, 242; spilota, 216, 258; Steinhelli, 240; Stevensi, 248; subsulcata, 221; suffusa, 229; tarsalis, 220; tetraspilota, 254; Theimei, 231; virescens, 223; virginella, 228; viridipennis, 234; viridi-pustulata, 226; vittata, 231; xanthoptera, 250; zonata, 239.
 Diadema, 96, 97, 98, 101, 102, 105, 106, 107, 108; setosum, 96, 113, 179, 201, 212; ambulacra of, 95.
 Diadematidae, 98, 112, 113, 179, 198, 201, 209; perignathic girdle in, 179, 198; structure of edges of plates at sutures, 98.
 Diaptomus castor, 296; orientalis, 296.
 Dicasticus, 327; celatus, 319, 329; laticollis, 319, 328; quadrimus, 319, 328, 336.
 Dicheilonema, 175.
 Difficulties against natural selection

- as a theory of the origin of species,
G. J. Romanes on, 338.
- Diodesma*, 61.
- Diphasia bipinnata*, 136, 159.
- Diplopoda*, 162, 163, 166, 167, 168,
172.
- Discoidea*, 207; *cylindrica*, 182.
- Distoma Boscii*, 177, 178; *clava*, 178;
incerta, 177, 178.
- Ditoma angustula*, 122; *rugicollis*, 117.
- Dog, brain of, 3, 4.
- Domestic dog, brain of, 3.
- Dorocidaris*, 179, 185, 186; *papillata*,
179, 181, 209.
- Duncan, Prof. P. M., on Anatomy of
Ambulacra of Recent Diadematidæ,
25; on *Perignathic Girdle of the*
Echinoidea 179.
- , and W. P. Sladen on test of
Arbaciadæ, 25.
- Dynamena tubulosa*, 137.
- Echinanthus rosaceus*, 179, 203, 204,
212.
- Echinarachnius*, 203; *mirabilis*, 207;
parma, 207.
- Echinidæ*, *perignathic girdle* in, 179,
194.
- Echinodiscus auritus*, 203, 207; *bi-*
foris, 207.
- Echinoidea*, Prof. P. M. Duncan on *peri-*
gnathic girdle of, 179.
- , 96, 110, 113, 179, 182, 184,
186, 187, 188, 193, 194, 200, 203,
207, 209.
- Echinometra*, 179; *lucunter*, 179,
200, 201; *subangularis*, 179, 201,
211.
- Echinometradæ*, 179, 198, 209; *peri-*
gnathic girdle in, 179, 198.
- Echinothrix*, 96; *calamaris*, 106; *De-*
sori, 101, 106, 114, 202, 212.
- Echinus*, 179, 180, 185, 197, 198, 199,
202, 208; *esculentus*, 179, 197;
norvegicus, 178, 194, 196, 210.
- Ectitheis*, 325; *divisus*, 319, 326, 336.
- Ectomicrus*, 59, 60, 129; *aper*, 129,
131; *pubens*, 60, 78, 80; *rugicollis*,
60, 78, 79, 129; *setosus*, 129.
- Ectozoa*, 178.
- Endophlæus*, 59, 60; *exsculptus*, 62;
serratus, 60, 61.
- Enhydra*, brain of, 20.
- Entomostraca*, coll. by Mr. A. Haly
in Ceylon, G. S. Brady on, 293.
- Entozoa*, 175, 260.
- Erotylathris*, 60, 127; *costatus*, 60,
75, 84, 127.
- Estheria Hislopi*, 294.
- Eucorybas crotalus*, 171.
- Eupleres*, brain of, 7.
- Eurycercus acanthocercoides*, 295.
- Evolution of species by independent
variation, G. J. Romanes on, 348.
- Felidæ*, brain of, 5.
- Filaria*, 175, 261; *horrida*, 175; *rheæ*,
175; *strongylina*, 261.
- Forcipules copultrices*, 164, 165,
172.
- Gadus*, worm in tail of, 187.
- Galerucinæ*, 213.
- Galictis*, brain of, 15.
- Galidia*, brain of, 8.
- Ganglion cells*, 170; of *Glomeris*, 168,
170; of *Sphærotherium*, 170.
- Gattya*, 155; *humilis*, 156, 161.
- Gempylodes*, 59, 60; *Lewisii*, 60, 73,
84.
- Genetta*, brain of, 6.
- Geotrupes stercorarius*, 87.
- Gervais, Prof., on brain of *Carni-*
vora, 1.
- Glomeridæ*, 162, 172.
- Glomeris*, 163, 165, 168, 171; *margi-*
nata, 171.
- Glyciphagus*, 269, 274; *dispar*, 271,
272, 273, 275, 280, 283, 284; *dispar*,
female of, 280; *dispar*, male of, 282;
hericius, 270; *ornatus*, 274; *palmi-*
mifer, 270; *platygaster*, 2 1, 272,
274, 280, 281, 282, 283, female of,
275, larva of, 280, male of, 278,
nymph of, 279; *plumiger*, 270.
- Glyphostomata*, 181, 187, 209.
- Glyptocryptus*, 58, 60; *brevicollis*, 60,
64.
- Goniocidaris geranioides*, 179, 185,
210.
- Grisonia*, brain of, 16.
- Gulo*, brain of, 17.
- Gyponychus*, 329.
- Halicornaria cornuta*, 153, 161; *mi-*
trata, 153, 160.
- Haly, A., *Entomostraca* coll. in Ceylon
by, 293.
- Harpacticidæ* (Ceylon), 296.
- Helictis*, brain of, 18.
- Herpestes*, brain of, 7, 8.
- Heteroplon*, 156.
- Honey-bee, antennæ of, 84.
- Hyæna*, brain of, 9.
- Icticyon venaticus*, brain of, 2, 4.
- Ictonyx*, brain of, 18.
- Ilyocryptus Hayli*, 294; *sordidus*,
295.
- Inutility of specific characters, 344.

- Ithris*, 59, 122; *decisa*, 123; *oculata*, 122; *sculpturata*, 60, 72.
Ithyporinæ (African), 319.
- Jackal, brain of, 3.
 Japanese *Colydiidæ*, list of, 60.
Julidæ, 166, 168.
Julus, 167, 168, 169.
- Krueg, on Carnivorous brain, 2.
- Labromimus*, 58, 60, 118; *variegatus*, 60, 65, 84.
Laganidæ, 207.
Laganum depressum, 179, 203, 206, 212.
 Langley, M., on brain of dog, 2.
Laparocerus, 327.
 Larva of *Glyciphagus platygaster*, 28.
Leptoglyphus, 59, 60, 125; *cristatus*, 125; *vittatus*, 60, 75.
Leptopinæ, African, 319.
Leptops, 329.
Lerneæ branchialis, 178.
 Leuret and Gratiolet, MM., on brains of Carnivora, 1.
Leydigia acanthocercoides, 295.
Limnadia Hislopi, 294.
Limnadiidæ, 294.
 Localities of new African *Cureulionidæ*, 318.
Lophyropoda, 297.
Lota, 178.
Loxococoncha alata, 312; *avellana*, 313; *elongata*, 313; *gibbera*, 312; *papillosa*, 313; *sagittalis*, 312.
Lutra, brain of, 20.
Lycaon pictus, brain of, 3, 4.
Lycetus, 117.
Lyncæidæ, 295.
Lyncodaphnidæ, 294.
Lytocarpus, 154; *myriophyllum*, 155; *ramosus*, 154, 161.
- Machlotes*, 76, 127, 128; *porcatus*, 76.
Macrocystis pyrifera, 133.
Macrothrix triserialis, 295.
Mecedanops ornamentalis, 123.
 Median suture of *ambulacra*, 105.
Melanthalia abscissa, 133.
 Meles, brain of, 14.
Mellita, 203; *testudinata*, 207.
Mellivora, brain of, 14.
Metopiestes erosus, 123; *tubulus*, 123, 131.
 Michael, A. D., on some undescribed Acari of genus *Glyciphagus* found in Moles' nests, 269.
Microcyphus zigzag, 179, 192, 210.
- Micropyga tuberculata*, 95, 110, 114.
Microvonus squalidus, 118, 131.
Minyad (*Minyas torpedo*) from N.W. Australia, by Prof. F. J. Bell, 114.
Minyas torpedo, 114, 116; zoological affinities of, 115.
 Mivart, St. George, on Cerebral Convulsions of Carnivora, 1.
Moina submucronata, 294.
 Morphology test of *Arbaciadæ*, Prof. Duncan and W. P. Sladen on, 25.
Mustela, brain of, 17.
Myrmica ruginodis, 88.
- Naidomorpha*, 267.
Nais, 264; *appendiculata*, 264, 266, 268; *escharosa*, 268; *lurida*, 264, 266, 267, 268; *serpentina*, 266.
Nandiniæ, brain of, 6.
Nasua, brain of, 11.
 Natural Selection not a theory of the origin of species, G. J. Romanes on, 345.
Nautactis, 116.
Neiphagus dentatus, 319, 333, 336; *fascicularis*, 319, 334.
Nematoideum Equi Caballi, 288.
Neoplatus, 119; *fasciatus*, 120.
Neotrichus hispidus, 60, 61, 84, 117; *serratus*, 117, 131.
Neurospongium, 169.
 New African genera and species of *Cureulionidæ*, by F. P. Pascoe, 318.
Normania avellana, 313.
 Notes on Entomostraca collected by Mr. A. Haly in Ceylon, by G. S. Brady, 293.
 Nymph of *Glyciphagus platygaster*, 279.
- Obelia longissima*, 136.
Oceanactis rhododactyla, 116.
Oligochaeta, 265.
Ophidonais and *Slavina*, by C. Bousfield, 264.
 — *serpentina*, 266, 268.
 Origin of Species, G. J. Romanes on, 337.
Ostra nodulosa, 319, 329.
Ostracoda (Ceylon), 297.
Otaria, brain of, 23.
Otiorthynchinæ (African), 319.
Otocyon megalotis, brain of, 3.
Oxylæmus, 74.
Oxyuris obesa, 176; *curvula*, 176.
- Pachylon Gorhami*, 79.
Pamphæa deficiens, 319, 322.
Paradoxostoma cingalense, 315.
Paradoxurus, brain of, 6.

- Parasites, notes on, by T. Spencer Cobbold, 174.
- Pascoe, F. P., on new African Genera and Species of Curculionidæ, 318.
- Penthelispa, 77; alternans, 128; crassicornis, 129; nitidicollis, 129.
- Perignathic girdle of Echinoidea, by Prof. P. M. Duncan, 179; terminology of parts of perignathic girdle, 181.
- Peripatus, 167, 170; capensis, 170.
- Peristhenes adnotus, 319, 335, 336.
- Peristomial region of ambulacra, 103.
- Philothermus, 59; depressus, 78.
- Phoca, brain of, 21.
- Phyllacanthus imperialis, 179, 186, 210.
- Phyllopoda, 294.
- Physiological Selection, an additional suggestion on the Origin of Species, Dr. G. J. Romanes on, 337; summary and conclusion, 397.
- — —, or Segregation of the Fit, 350; argument from divergence of character, 385; argument from geographical distribution, 388; argument from inutility of specific differences, 382, arguments on physiological selection, 354, 365; argument from prevention of intercrossing, 378; argument from sterility between species, 366; general summary and conclusion, 397.
- Piazonias peregrinus, 318, 319; macer, 318, 320; velatus, 319.
- Pinnipedia, brain of, 21, 24.
- Platanista gangetica, 177.
- Platyomicus aridus, 326, 336; cordipennis, 326; echinus, 326.
- Plotactis, 115, 116.
- Plumularia lagenifera, 157, 158, 161; multinoda, 157, 161.
- Plumularidæ, 155.
- Pontocypris Davidsoni, 304; nitida, 303; trigonella, 304.
- Procyon, brain of, 10.
- Prolyctes bituberculatus, 128, 131.
- Proteles, brain of, 9, 10.
- Prototracheata, 167.
- Psalistus, 335.
- Psammechinus miliaris, 179, 197, 211.
- Pseudotarphius, 58; Lewisii, 60, 63.
- Pynomerini, 60.
- Pynomerus, 59; alternans, 128, 131; crassicornis, 129; distans, 129, 131; nitidicollis, 129; sculpturatus, 60, 77; vilis, 60, 77.
- Races in regard to physiological selection, 358.
- Relations between survival of fittest and segregation of fit, 395.
- Rhea, 174, 175.
- Rhizocephala, 262.
- Romanes, G. J., on Physiological Selection, 337-411.
- Sacculina, 261.
- Salmacis, 179, 187, 191, 192, 196; bicolor, 179, 187, 210.
- Saphicus variegatus, 319, 336.
- Sargassum, 138, 139, 142, 159.
- Sclerostoma struthionis, 263.
- Scobius, 326.
- Scolopendra, 167.
- Seals, brains of, 21, 23.
- Selaginopsis, 145.
- Sense-cells and ganglion-cells in Sphærotherium, 170.
- Sertularia, 138, 139; ampletens, 141, 142, 159; aperta, 138, 150, 159; bispinosa, 140; crinis, 139, 140, 159; crinoidea, 141, 159; elongata, 140, 159; geminata, 143; humilis, 143; megalocarpa, 142, 143, 159; minima, 138, 139, 159; operculata, 138, 148; unguiculata, 144; unilateralis, 139, 159.
- Sertularella, 133; capillaris, 133, 158; crassipes, 133, 158; cuneata, 134, 158; diffusa, 136, 158; Johnsoni, 133; limbata, 134, 158; margaritacea, 133, 158; trimucronata, 135, 158; trochocarpa, 135, 158.
- Sharp, D., on Colydiidæ from Ceylon, 117; on Colydiidæ from Japan, 58.
- Sideroactylus delectans, 318, 320; Oberthuerii, 318, 320; puellaris, 318, 320.
- Simondsia paradoxa, from Hog, 261.
- Sladen, W. P., and Prof. Duncan, on test of Arbaciadæ, 25.
- Slavina and Ophidonais, by E. C. Bousfield, 264.
- — —, 264; appendiculata, 264, 268; lurida, 267, 268; serpentina, 267, 268.
- Species in regard to Physiological Selection, 361.
- Sphadasmus, 335.
- Sphærotherium, anatomy of, G. C. Bourne on, 161; copulatory appendages of, 164; stridulating organ of, 165, 172; tergites of body, 162; trachea, primitive charac. of, 167; tracheal plates, 163.
- obtusum, 162, 164, 172, 173; retusum, 162, 164, 172, 173.
- Sphærolaria, 261.

- Spirobolus, 169.
 Spiroptera megastoma, 261; strongylina, 261.
 Stenophida linearis, 319, 336.
 Sterility between species, G. J. Romanes on, 340.
 Stiamus brachyurus, 319, 325, 336.
 Stigmatrachelus alternans, 322; flexuosus, 319, 324; longiceps, 319, 322; ruptus, 319, 322; vittatus, 319, 322.
 Straticus, 326; funestus, 319, 327.
 Stridulating-organ of Sphærotherium, 165, 172.
 Strongylocentrotus, 96, 101, 105, 110, 179, 198, 200; lividus, 179; perignathic girdle in, 198, 211.
 Strongylus Arnfeldi, anatomy of, 285.
 — Axei (Cobb.), preceded by remarks on its affinities, by Dr. T. Spencer Cobbold, 259.
 —, 260; armatus, 288; Arnfeldi, 285; Axei, 262, 263, 264; contortus, 260; Douglassii, 260, 262, 263; filaria, 285; micrurus, 285; pergracilis, 260; tetracanthus, 286.
 — tetracanthus, structure and development of, 286, 288, 291; cocoon-forming habit of, 290; literature concerning, 291.
 Structure of edges of plates at sutures in Diadematidæ, 98.
 Struthio molybdophanes, Strongyloid worm in, 263.
 Suricata, brain of, 8.
 Swamping effects of intercrossing, 343.
 Sympanotus, 58; pictus, 60, 62, 84.
 Synchitini, 60.
 Synthecium elegans, 137, 138; ramosum, 137, 159.
 Systates laticollis, 319, 327.
 Tanyrhynchinæ (African), 319.
 Tanyrhynchus ellipticus, 319, 330, 336.
 Tarphiosoma echinatum, 119; fasciata, 120; luridum, 119.
 Tarphius pilosus, 119.
 Temnopleuridæ, 179, 187, 192, 193, 195, 200, 208, 209; perignathic girdle in, 187.
 Temnopleurus toreumaticus, 179, 190.
 Tere dolæmus, 59, 124; biplagiatus, 124; guttatus, 60, 74; politus, 60, 74, 84, 124; similis, 124.
 Tere dus, 60, 74; biplagiatus, 124; politus, 60.
 Tetrapygus nigra, 57.
 Thecocladium, 149; flabellum, 149, 160.
 Thuiaria, 136, 144, 145, 149; diaphana, 145, 160; heteromorpha, 147, 160; Hippisleyana, 146, 160; interrupta, 145, 159; polymorpha, 149; ramosissima, 146, 160.
 Thyroderus, 58, 60; porcatus, 60, 83, 84.
 Timola, 331; suturalis, 319, 332.
 Tracheal system of Sphærotherium, 166.
 Trachypholis, 120; Erichsoni, 120, 131; fasciculata, 120.
 Trichodactylus, 274; anonymus, 274.
 Trichonema arcuata, 288.
 — stage of growth of Strongylus tetracanthus, 291.
 Trionus, 59; opacus, 60, 70, 122, 131.
 Tyroglyphidæ, 270.
 Tyroglyphus, 270; anonymus, 274; longior, 274; setiferus, 273.
 Ursus, brain of, 19.
 Vespa Crabro, antennary organs of, 170.
 Viverra, brain of, 5, 10.
 Viverrinæ, brain of, 24.
 Watson, Rev. R. Boog, on Cerithiopsides from N. Atlantic and Madeira, 89.
 Wilder, Prof. Burt, on brains of dog and cat, 2.
 Wolf, brain of, 3.
 Worm in tail of Gadus, 178.
 — from the mouth of a snake, 177.
 Xestoleberis curta, 314; intermedia, 314; sulcata, 314, tumifacta, 314; variegata, 314.
 Xuthia, 58, 59; niponica, 60, 71, 123; parallela, 60, 70, 122, 131.
 Xylophagous insects of Japan, 58.
 Zephronia, 162, 166.
 Zygotinæ (African), 319.

END OF THE NINETEENTH VOLUME.